











- [14] P. Shah and Y. Luximon, "Three-dimensional human head modelling: a systematic review", in *Theoretical Issues in Ergonomics Science*, 2018. DOI: 10.1080/1463922X.2018.1432715Y.
- [15] K.M. Robinette et al., "The CAESAR project: a 3-D surface anthropometry survey". in *Proc. of Second International Conference on 3-D Digital Imaging and Modeling*, Ontario, Canada, 1999, pp. 380-386, <https://doi.org/10.1109/IM.1999.805368>
- [16] T. Perret-Ellena et al., "3D Anthropometric investigation of Head and Face characteristics of Australian Cyclists", in *Procedia Engineering*, 112, 2015, pp. 98-103, <https://doi.org/10.1016/j.proeng.2015.07.182>
- [17] H.P. Lee et al., "Development of an Anthropometric Database Representing the Singapore Population", in *Proc. of 6th International Conference on 3D Body Scanning Technologies*, Lugano, Switzerland, 2015, pp. 27-28, <http://dx.doi.org/10.15221/15.234>
- [18] H. S. Baik et al., "Facial soft-tissue analysis of Korean adults with normal occlusion using a 3-dimensional laser scanner", in *American journal of orthodontics and dentofacial orthopedics*, 131(6), 2007, pp. 759-766, <https://doi.org/10.1016/j.ajodo.2005.08.038>
- [19] D.S. Ma et al., "The Chicago face database: A free stimulus set of faces and norming data", in *Behavior research methods*, 47(4), 2015, pp. 1122-1135. <https://doi.org/10.3758/s13428-014-0532-5>
- [20] L. Talbert et al., "A 3D analysis of Caucasian and African American facial morphologies in a US population", in *Journal of orthodontics*, 41(1), 2014, pp.19-29. <https://doi.org/10.1179/1465313313Y.0000000077>
- [21] R. Ball et al., "A comparison between Chinese and Caucasian head shapes", in *Applied Ergonomics*, 41, no. 6, 2010, pp. 832-839. <https://doi.org/10.1016/j.apergo.2010.02.002>